

# A search for hidden white dwarfs in the *ROSAT* EUV survey – II. Discovery of a distant DA+F6/7V binary system in a direction of low-density neutral hydrogen

M. R. Burleigh,<sup>1</sup> M. A. Barstow<sup>1</sup> and J. B. Holberg<sup>2</sup>

<sup>1</sup>*Department of Physics and Astronomy, University of Leicester, University Road, Leicester LE1 7RH*

<sup>2</sup>*Lunar and Planetary Laboratory, Gould-Simpson Building, University of Arizona, Tucson AZ 85721, USA*

Accepted 1998 June 12. Received 1998 June 4; in original form 1997 October 20

## ABSTRACT

The *ROSAT* Wide Field Camera (WFC) survey of the extreme ultraviolet (EUV) has provided us with evidence for the existence of a previously unidentified sample of hot white dwarfs in unresolved, detached binary systems. These stars are invisible at optical wavelengths due to the close proximity of their much more luminous companions (spectral type K or earlier). However, for companions of spectral type  $\sim$ A5 or later the white dwarfs are easily visible at far-ultraviolet wavelengths, and can be identified in spectra taken by *IUE*. 16 such systems have been discovered in this way through *ROSAT*, *EUVE* and *IUE* observations, including four identified by us in Paper I. In the present paper we report the results of our continuing search during the final year of *IUE* operations. One new system, RE J0500–364 (DA+F6/7V), has been identified. This star appears to lie at a distance of  $\sim$ 500–1000 pc, making it one of the most distant white dwarfs, if not the most distant, to be detected in the EUV surveys. The very low line-of-sight neutral hydrogen volume density to this object could place a lower limit on the length of the  $\beta$  CMa interstellar tunnel of diffuse gas, which stretches away from the Local Bubble in a similar direction to RE J0500–364.

In this paper we also analyse a number of the stars observed where no white dwarf companion was found. Some of these objects show evidence for chromospheric and coronal activity. Finally, we present an analysis of the previously known WD+active F6V binary HD 27483 (Böhm-Vitense 1993), and show that, at  $T \approx 22\,000$  K, the white dwarf may be contributing significantly to the observed EUV flux. If so, it is one of the coolest such stars to be detected in the EUV surveys.

**Key words:** binaries: general – white dwarfs – ISM: general – ultraviolet: stars – X-rays: stars.

## 1 INTRODUCTION

The vast majority of the >2000 known white dwarfs (McCook & Sion, in preparation) are isolated stars discovered at optical wavelengths by virtue of their photometric colours or proper motions. In either case, there is a strong bias against detecting any white dwarfs in unresolved binary systems. A companion star of type K or earlier will completely dominate the optical spectrum of the white dwarf, effectively rendering it invisible. Indeed, Sirius B, the first white dwarf to be discovered, would never have been resolved from the A1 dwarf Sirius were it not for the close proximity of the system to Earth (2.64 pc).

Prior to the *ROSAT* and *Extreme Ultraviolet Explorer* (*EUVE*) surveys, a small number of unresolved white dwarf/main sequence binaries had been discovered serendipitously. For example, the white dwarf in the V471 Tauri system was found as a result of an

eclipse by its active K2V companion. In addition, a number of white dwarfs have been accidentally discovered during various far-ultraviolet (far-UV) observations of the normal stellar companions by the *International Ultraviolet Explorer* (*IUE*), e.g.,  $\zeta$  Cap (Böhm-Vitense 1980), 56 Peg (Schindler et al. 1982) and 4  $\sigma^1$  Ori (Johnson & Ake 1986), although Shipman & Geczi (1989) systematically studied the then existing *IUE* archive for white dwarf companions to G, K and M stars, and found no further examples.

Now, however, the extreme-ultraviolet (EUV) surveys of the *ROSAT* Wide Field Camera (WFC; Pounds et al. 1993; Pye et al. 1995) and *EUVE* (Bowyer et al. 1994, 1996) have provided evidence for the existence of a substantial sample of these previously unknown white dwarfs, through the detection of EUV radiation. 16 new systems have been discovered in this way, including  $\beta$  Crateris (A2IV+WD; Fleming et al. 1991), KW Aur C (F4V+DA; Hodgkin et al. 1993), HD 18131 (K0IV+DA; Vennes

**Table 1.** *ROSAT* WFC/PSPC and *EUV*E count rates ( $1000 \text{ s}^{-1}$ ).

<i>ROSAT</i> No.	Name <sup>†</sup>	WFC		PSPC		<i>EUV</i> E	
		S1	S2	(0.1–0.4 keV)	(0.4–2.4 keV)	100 Å	200 Å
WD Binaries							
RE J0500–36 <sup>a</sup>		$20 \pm 5$	$47 \pm 8$	no det.	no det.	$21 \pm 6$	no det.
RE J0420+13 <sup>b</sup>	HD 27483	$15 \pm 4$	$27^{\dagger}$	$98 \pm 14$	$33 \pm 9$	no det.	no det.
Non-WD Systems							
2RE J0222+50	BD+49° 646	$14 \pm 3$	$17 \pm 5$	$258 \pm 20$	$195 \pm 18$	no det.	no det.
2RE J0232–02		$15^{\dagger}$	$45 \pm 11$	no det.	no det.	no det.	no det.
2RE J0530–19	SAO 150508	$8 \pm 3$	$16 \pm 5$	$159 \pm 18$	$195 \pm 20$	no det.	no det.
2RE J0534–15	HD 36869	$8 \pm 3$	$19 \pm 5$	$221 \pm 22$	$217 \pm 22$	no det.	no det.
2RE J0544–22	GI 216B	$7 \pm 3$	$22 \pm 5$	$261 \pm 23$	$82 \pm 13$	no det.	no det.
RE J0613–23	HR 2225	$18 \pm 3$	$35 \pm 9$	$517 \pm 30$	$256 \pm 21$	no det.	no det.
2RE J0640–03	HD 295290	$9 \pm 3$	$32 \pm 8$	$304 \pm 28$	$281 \pm 27$	no det.	no det.
2RE J0710+45	HD 54402	$8 \pm 3$	$26 \pm 7$	no det.	no det.	no det.	no det.
2RE J0813–07	SAO 135659	$14 \pm 6$	$55 \pm 16$	$207 \pm 35$	$195 \pm 34$	no det.	no det.
RE J0823–25 <sup>c</sup>		$52 \pm 7$	$83 \pm 9$	$175 \pm 24$	no det.	$28 \pm 11$	no det.
2RE J1212+77	HR 4646	$13^{\dagger}$	$18 \pm 5$	$157 \pm 42$	$114 \pm 36$	no det.	no det.

<sup>†</sup> upper limit<sup>a</sup> Detected in the First *EUV*E Source Catalog only (Bowyer et al., 1994)<sup>b</sup> *ROSAT* WFC Bright Source Catalogue detection only (Pounds et al., 1993)<sup>c</sup> HD 70907 and a  $V \approx 11$  companion were both observed as possible counterparts to this source

et al. 1995), RE J0357+283 (K2V+DA; Jeffries, Burleigh & Robb 1996), and the latest two, RE J0702+129 (K0Ve+DA; Vennes et al. 1997b), and HR 2875 (B5V+WD; Vennes, Berghoefer & Christian 1997a; Burleigh & Barstow 1998). Detailed studies of groups of these objects have been undertaken by Barstow et al. (1994) and Burleigh, Barstow & Fleming (1997, hereafter Paper I). Positive identifications have been made in each case through follow-up observations in the far-UV with *IUE*, since for companions later than  $\sim A5$  the hot white dwarf is easily visible at these wavelengths (the companion to the B5V star HR 2875 was identified through an *EUV*E spectrum, since the B star still dominates the spectrum at far-UV wavelengths).

It is well established that over half of all stars are members of binary or multiple systems, yet the overwhelming majority of catalogued white dwarfs are isolated objects. The new population of optically hidden white dwarfs emerging from the EUV surveys has profound implications, therefore, for our knowledge of the white dwarf luminosity function, space density and formation rate (e.g. Fleming, Liebert & Green 1986). Observations of white dwarfs in detached binary systems also allow us to place constraints on their evolutionary models (e.g. de Kool & Ritter 1993).

In this paper we report the results of a continuing search for more of these binaries in the *ROSAT* WFC catalogue, during the final year of *IUE* operations. We have discovered one new unresolved white dwarf+main sequence star binary (RE J0500–362).<sup>1</sup> In the cases where no white dwarf was detected, a number of the target stars show evidence for chromospheric activity. We also present an analysis of the previously known white dwarf+active star system HD 27483 (Böhm-Vitense 1993). We show that the white dwarf is contributing to the EUV flux, and should be included in the growing list of EUV-emitting, optically hidden hot white dwarfs in binary systems.

## 2 DETECTION OF THE SOURCES IN THE *ROSAT* WFC EUV SURVEY

The *ROSAT* WFC EUV all-sky survey was conducted between 1990 July and 1991 January. Two broad-band filters were utilized (designated S1 and S2), and most of the count rates quoted in this paper (see Table 1) are taken from the 2RE catalogue (Pye et al. 1995). This revised list contains 479 EUV sources, as compared with 383 in the original Bright Source Catalogue (Pounds et al. 1993). The 2RE catalogue was compiled from the original survey data with improved methods for source detection, background screening, etc. The resulting count rates are equivalent to on-axis, at-launch values.

The *ROSAT* PSPC X-ray survey was conducted simultaneously with the WFC. The soft (0.1–0.4 keV) and hard (0.4–2.4 keV) band count rates (Table 1) were obtained via the World Wide Web from the on-line All-Sky Survey Bright Source Catalogue, maintained by the Max-Planck Institute in Germany (Voges et al. 1996). All of the X-ray flux from hot white dwarfs is expected to lie within the soft band.

The *EUV*E all-sky survey was conducted in four passbands between 1992 July and 1993 January. These count rates are also given in Table 1, and are mainly taken from the Second *EUV*E Source Catalog (Bowyer et al. 1996) which, like the *ROSAT* WFC 2RE catalogue, includes better source detection algorithms and improved reliability.

### 2.1 Selection of candidate hidden white dwarfs

Hot white dwarfs in unresolved binaries with companions of spectral type K or earlier are virtually impossible to discern from optical spectra alone. However, it is possible to identify these hidden white dwarfs unambiguously in far-UV spectra taken by *IUE*. The problem is how to select likely candidates from just their EUV and soft X-ray count rates.

Most of the hot white dwarfs discovered by *ROSAT* have very soft spectra compared to normal stars, particularly where the interstellar

<sup>1</sup> Originally reported, with a preliminary analysis, in the proceedings of the 1996 European White Dwarf Workshop by Burleigh & Barstow (1997).

hydrogen column density is low. The ratio of the WFC S2 to S1 count rates can often exceed a factor of 2. Additionally, no photons are usually detected from a white dwarf above the 0.28-keV carbon edge of the ROSAT PSPC. All other X-ray and EUV-emitting astronomical objects generally have spectra extending to higher energies. Thus many of the stars originally selected for observation by *IUE* were relatively bright EUV sources with very similar colours to known hot white dwarfs (e.g., KW Aur C; Hodgkin et al. 1993). The success rates of the early searches by, for example, Barstow et al. (1994) were very high, and appeared to represent the tip of an iceberg.

However, many of the white dwarfs in the ROSAT WFC survey are relatively faint EUV sources, indistinguishable by count rate ratios alone from coronally active objects. Further selection criteria, in addition to simple EUV colour and brightness, needed to be applied. In Paper I we conducted a search for fainter, less obvious examples of these binary systems, with a  $\sim 40$  per cent success rate. Candidates were selected, in particular, from normal stars that had been observed in the WFC optical identification programme (Mason et al. 1995), and where little or no evidence of activity had been found. In some cases, e.g., HD 2133 (RE J0024–741), a hidden white dwarf was indeed detected in an *IUE* SWP spectrum. In others, e.g., HR 2468 (RE J0637–613), the *IUE* observations revealed evidence of chromospheric and coronal emission that had eluded the optical identification team.

In this paper we report the results of a further search for hidden white dwarfs during the last year of *IUE* operations (1995/96). 13 candidates were selected and observed, including many that were detected for the first time in the reprocessed ROSAT WFC 2RE data.

These are, in general, faint EUV sources. In most cases, the target stars were not known to be active, and had S2/S1 count rate ratios  $> 2$ . In a separate optical programme, a number of the unidentified 2RE fields were observed to try to determine the counterpart to the EUV source. In some cases, e.g., RE J0500–364, 2RE J0222+503 and 2RE J0232–025, only one relatively faint star ( $V > 10$ ) was visible in the field. Although a few EUV sources (mainly red dwarfs) are active stars with  $V > 10$  (e.g., Proxima Cen, M5Ve,  $V = 11.1$ ), most are much brighter than this. Thus, for these sources, a hidden hot white dwarf companion was a feasible explanation for the EUV radiation, and they were added to the *IUE* target list.

## 2.2 UV spectroscopy

A log of all the far-UV observations is given in Table 2. The white dwarf companion to HD 27483 was serendipitously discovered in 1992 October by Böhm-Vitense (1993), independently of the ROSAT survey, during an *IUE* survey of Hyades F stars. RE J0500–364 was first observed in a short half-hour exposure in 1995 November (SWP56217), and a faint white dwarf was detected. A follow-up spectrum, with an 8-h exposure to achieve the required signal-to-noise ratio, was obtained a month later.

The programme was unfortunately cut short in 1996 February when a gyro failed on *IUE*, limiting observations to objects for which a bright ( $V < 8$ ) guide star was available. This was not the case for any of our remaining targets.

Two spectra were affected by the so-called ‘159DN anomaly’ (see Table 2). A number of pixels were assigned the flux value

**Table 2.** Log of *IUE* observations.

Name	SWP No.	LWP No.	Date	Exp. (s)	Observer	notes
WD Binaries						
RE J0500–364	56217		1995/323	1800	SO	
	56338		1995/358	29400	SO	159DN problem
		31729	1995/323	1800	SO	
HD 27483	45940		1992/273	2100	Böhm-Vitense	
Non-WD systems						
2RE J0222+50	56333		1995/358	1800	SO	
2RE J0232–02	56272		1995/340	1800	SO	
		31800	1995/340	1800	SO	
SAO 150508	56195		1995/317	1800	SO	
		31700	1995/317	1200	SO	
HD 36869	56196		1995/317	1800	SO	
		31701	1995/317	360	SO	
Gl 216B	56194		1995/317	1800	SO	
		31699	1995/317	120	SO	
HR 2225	56206		1995/319	1800	SO	
		31715	1995/319	60	SO	
HD 295290	56211		1995/322	1800	SO	
		31726	1995/322	1200	SO	159DN problem
HD 54402	56193		1995/317	1800	SO	
		31698	1995/317	300	SO	
SAO 135659	56273		1995/340	1800	SO	
		31801	1995/340	1200	SO	
HD 70907	56344		1995/359	1800	SO	
		31836	1995/359	300	SO	
RE J0823–25 <sup>†</sup>	56266		1995/338	1800	SO	
		31796	1995/338	300	SO	
HR 4646	56393		1996/009	100	SO	

SO = Service Observation

<sup>†</sup>  $V \approx 11$  companion to HD 70907

**Table 3.** Physical parameters of the companion stars in the WD binaries.

RE No.	Cat. Name	SpT	V	$d$ est. (pc)	references
RE J0500–36		F6/7V	13.7	520–1250	
RE J0420+13	HD 27483	F6V	6.17	47.6	Böhm-Vitense (1993)

**Table 4.** Physical parameters of the stars observed where no white dwarf was detected.

RE No.	Cat. Name	Spectral type	V magnitude	references
2RE J0222+50	BD+49°646	?	10.1	SIMBAD
2RE J0232–02		K	14.8	SIMBAD
2RE J0530–19	SAO 150508	G5V	9.0	SIMBAD
2RE J0534–15	HD 36869	G2V	8.0	SIMBAD
2RE J0544–22	Gl 216B	K2V	6.15	SIMBAD
RE J0613–23	HR 2225	G5V	6.39	SIMBAD
2RE J0640–03	HD 295290	G0	9.1	SIMBAD
2RE J0710+45	HD 54402	K0	7.7	SIMBAD
2RE J0813–07	SAO 135659	K0	8.8	SIMBAD
RE J0823–25	HD 70907	F3IV/V	8.8	SIMBAD
		late	11	Mason et al. (1995)
2RE J1212+77	HR 4646	A5m	5.14	SIMBAD

**Table 5.** Temperatures and gravities for the white dwarfs from homogeneous model fits.

Binary	$\log g$	Temp K	90 per cent error K	Mass $M_{\odot}$	Radius $R_{\odot}$	$d_{\text{wd}}$ (pc)	Estimated $V$
RE J0500–36	7.0	36,800	33,900–41,000	0.38	0.032	1050	17.8
	7.25	38,260	35,350–42,900	0.40	0.025	830	17.9
	7.5	42,000	36,400–45,700	0.47	0.020	740	17.9
	8.0	45,000	40,500–50,500	0.68	0.014	520	17.9
	8.5	50,500	45,400–57,700	0.97	0.009	390	18.0
HD 27483	9.0	60,500	51,900–67,500	1.21	0.006	270	18.1
	7.0	20,000	upper limit	0.37	0.032	103	13.9
	7.5	20,000	upper limit	0.40	0.019	61	13.9
	8.0	21,410	21,215–21,630	0.63	0.013	50	14.2
	8.5	22,000	21,910–22,780	0.94	0.009	44	14.5
	9.0	24,800	24,570–25,060	1.20	0.006	30	14.6

159DN when they should in fact have had a higher value. However, on close inspection it was found that only a small fraction of each of these two spectra was contaminated, and their scientific usefulness was not compromised.

### 3 IUE DATA REDUCTION

All of the spectra have been processed with NEWSIPS (New Spectral Image Processing System). NEWSIPS spectra contain a number of significant geometrical and photometric corrections which enhance the spectral signal-to-noise ratio and improve the photometric reliability of the data (e.g., a correction for the degradation of the detector with time; Bohlin & Grillmair 1988). Recent analysis by Garhart (1997) shows that, since 1993, the SWP camera sensitivity has in fact degraded at a rate greater than predicted, so that the current NEWSIPS calibration underestimates the SWP fluxes by up to 5 per cent. Archival IUE data from 1993–96 are now being reprocessed to include a new degradation correction. However, these data were not available to us when we were preparing this paper, although we would expect the effect of the new calibration to be relatively minimal on the determination of any white dwarf parameters (for example, the errors on the flux values for the RE J0500–364 IUE SWP data are typically of order  $\sim 10$  per cent anyway).

## 4 ANALYSIS

The method used to analyse the far-UV and EUV data for the hidden white dwarfs and the active stars detected and observed during this search has been discussed in detail in Paper I. A summary is given here.

### 4.1 Hidden white dwarfs – far-UV data

In binary systems like these, it is not possible to use the Balmer line profiles to measure temperature and gravity. However, the IUE SWP spectra can be used to estimate these parameters by matching the observed Lyman  $\alpha$  profile and the UV continuum (the region uncontaminated by the companion) with synthetic spectra. Unfortunately, it is not possible to get an unambiguous determination of  $T$  and  $\log g$  from a single spectral line. Instead, a range of possible models is determined by stepping through values of the surface gravity, from  $\log g = 7.0$  to  $9.0$ , and finding the best-fitting temperature and normalization at each point.

We compare the observed far-UV data with fully line-blanketed, homogeneously mixed H+He, LTE model atmospheres, spanning a temperature range from 20 000 to 100 000 K, and kindly supplied by Detlev Koester (e.g. Koester 1991). In this analysis we assume the white dwarfs have pure hydrogen envelopes. The spectral fitting is conducted with the XSPEC program (Schafer et al. 1991), which

calculates a chi-squared statistic for the fit between the data and the model, and which is then minimized by incremental steps in the free parameters. There is no need to take into account any interstellar component in the analysis of the Lyman  $\alpha$  profile, because for columns greater than a few  $\times 10^{19}$  the white dwarf is unlikely to be detected in EUV surveys.

The white dwarf radius and mass are calculated using the temperature-dependent evolutionary models of Wood (1995), which assume thick ( $10^{-4} M_{\odot}$ ) H layers, and the radii are then used to estimate the distance from the model normalization parameter [which is actually the solid angle of the star, equivalent to  $(\text{radius}/\text{distance})^2$ ]. The distances to the primaries in each case can be calculated from the spectral type and  $V$  magnitude, and are given in Table 3. The range of white dwarf temperatures and gravities that give the best match to the distance of the primary can then be estimated. The  $V$  magnitude of each white dwarf is estimated from the model flux at 5500 Å, and is given in Table 5.

#### 4.2 Hidden white dwarfs – ROSAT data

Once the temperature and gravity of each white dwarf has been estimated, the ROSAT EUV and soft X-ray fluxes can give an indication of the level of photospheric opacity in the stellar atmosphere, by comparing them with predicted values for a pure H model. The ROSAT data are fitted independently from the IUE data, since contamination from elements heavier than H and He has an effect only at EUV and soft X-ray wavelengths. We fit the data from the two WFC filters, and the integrated count rate in the 0.1–0.28 keV PSPC band, within which all the white dwarf soft X-ray flux is expected to lie. It is possible that some EUV and X-ray emission might also originate from the normal stars in each system. Any detected PSPC flux at energies above 0.4 keV is an indication of the presence of an active companion (Table 6).

Once again we utilize Detlev Koester’s fully line-blanketed H+He models. These assume a homogenous distribution of hydrogen and helium, under LTE conditions, in the range  $-8 < \log \text{He}/\text{H} < -3$ . The temperature, gravity and normalization,

estimated from the fit to the IUE data, are frozen during the modelling, but the He/H ratio is allowed to vary. The interstellar H I column density is also estimated by assuming that the local ISM is not highly ionized (i.e., there is minimal He II absorption) and that the He I/H ratio is cosmic (0.1).

The fitting is again conducted using the XSPEC program. We consider a good fit to the data to correspond to the probability that a particular value of the reduced  $\chi^2$  ( $\chi_r^2$ ) can occur by chance to be 0.1 or greater (i.e., 90 per cent confidence), and a bad fit 0.01 or less (99 per cent confidence). The fits in between might not be very good, but cannot be ruled out with high confidence. In this analysis a good fit requires  $\chi_r^2 < 2.71$ , but until it exceeds 6.63 a model cannot be excluded with any certainty. Consequently, we list all model fits to the data for which  $\chi_r^2 < 6.63$  (Table 6).

#### 4.3 Non-detections and active stars

A number of the stars observed with IUE where no white dwarf was detected are probably coronally and chromospherically active. Some of these stars are also comparatively hard X-ray sources (i.e., they are detected in the 0.4–2.4 keV band of the ROSAT PSPC), and in some of the IUE LWP spectra chromospheric Mg II emission was seen at 2798 Å.

In these cases, estimates of the  $L_{\text{EUV}}/L_{\text{bol}}$  and  $L_{\text{x}}/L_{\text{bol}}$  ratios have been made as an indicator of the level of activity (Table 7). We adopt the methods outlined by Jeffries (1995) and Fleming et al. (1995) to estimate these parameters.

Where chromospheric Mg II emission was seen the line fluxes above the continuum level were measured using a simple Gaussian profile, fitted to each line, after the continuum had first been subtracted (the continuum flux was represented by a low-degree polynomial). Note that in each case the Mg II emission actually fills in an underlying absorption dip. However, it is impossible to estimate the depth of this absorption line, and thus the measurements presented here are only lower limits. The measured Mg II fluxes are listed in Table 7, and have been used to estimate  $L_{\text{MgII}}/L_{\text{bol}}$ .

**Table 6.** Column densities from homogeneous models for RE J0500–362 (assuming pure H composition).

Name	$\log g$	$T$	H I column	90 per cent error	Comment
RE J0500–364	7.0	36 800	$3.7 \times 10^{18}$	$1.7\text{--}7.0 \times 10^{18}$	
	7.25	38 260	$7.5 \times 10^{18}$	$5.0 \times 10^{18}\text{--}1.1 \times 10^{19}$	
	7.5	42 000	$1.7 \times 10^{19}$	$1.4\text{--}2.1 \times 10^{19}$	
	8.0	45 000	$2.4 \times 10^{19}$	$2.1\text{--}2.9 \times 10^{19}$	
	8.5	50 500	–	–	No fit
	9.0	60 500	–	–	No fit

No fit could be obtained to the ROSAT data for HD 27483.

**Table 7.** EUV and X-ray luminosities for the probable and confirmed active stars.

Name	$d$ (pc)	$L_{\text{EUV}}$ $\times 10^{29}$ ergs	$L_{\text{EUV}}/L_{\text{bol}}$ $\times 10^{-4}$	$L_{\text{x}}$ $\times 10^{29}$ ergs	$L_{\text{x}}/L_{\text{bol}}$ $\times 10^{-4}$	$f_{\text{MgII}}$ $\times 10^{-12}$	$L_{\text{MgII}}/L_{\text{bol}}$ $\times 10^{-4}$
SAO 150508	59	10.0	3.6	1.3	4.7	–	–
HD 36869	48	7.9	1.7	10.0	2.2	–	–
GI 216A	8	0.3	0.04	0.2	0.02	–	–
GI 216B	8	0.3	0.3	0.2	0.2	4.2	0.5
HR 2225	18	2.0	0.7	2.0	0.7	–	–
HD 295290	95	51.8	8.2	51.5	8.2	0.6	1.0
SAO 135659	34	11.4	0.7	4.5	0.4	1.0	1.1

## 5 DISCUSSION

### 5.1 Hidden white dwarf binaries


#### 5.1.1 RE J0500–364

The field of this EUV source (see Fig. 1) was surveyed by Mason et al. (1995) during the WFC optical identification programme. No evidence of activity was found in the cores of Ca II H & K in the spectrum of a  $\sim 13$ th magnitude star located at the centre of the source error box. Various stars were also examined outside the error box, but none could plausibly account for the EUV emission. The field was also examined on 1995 September 29/30 with the 2.3-m Steward Observatory telescope on Kitt Peak, as part of a programme to search for the optical counterparts to unidentified sources in the *ROSAT* WFC 2RE catalogue. A spectrum (Fig. 2) of the central star in the source error box was obtained with the Boller & Chivens Spectrometer and  $800 \times 1200$  blue-sensitive CCD. A 2.5-arcsec slit and  $600 \text{ line mm}^{-1}$  grating blazed at  $3658 \text{ \AA}$  were used, and the data were reduced with standard IRAF routines. Again, there was no evidence for activity. One untested proposition was that this central object could be hiding a hot white dwarf companion. The star was therefore added to our *IUE* target list for observations during 1995/96, and a faint,  $\approx 18$ th magnitude hot white dwarf companion was discovered (Fig. 3).


By comparing the relative line strengths and widths of the primary's optical spectrum with spectra in the atlas of Jacoby, Hunter & Christian (1984), we conclude that it most closely resembles an F6/7V star. It should be noted that the spectrum appears to be deficient in flux at the blue end, below  $\sim 4500 \text{ \AA}$ . As the star was observed at a relatively high airmass (sec  $Z = 2.94$ ), this deficiency in counts was probably caused by differential atmospheric dispersion. The spectral identification is strengthened when the *IUE* LWP spectrum (LWP31729, Fig. 4) is compared to spectra in the *IUE* Spectral Atlas (Wu et al. 1992). The data are very noisy, and have therefore been binned, but they most closely match stars in the range F5V–G0V. There also appears to be a slight excess of flux in the *IUE* SWP spectrum, at wavelengths  $> 1850 \text{ \AA}$ , above the level expected from the white dwarf (Fig. 3). If the excess is real, then it must be attributable to the primary. As  $\sim 13$ th magnitude mid-late G and K stars are not bright enough to be detected by *IUE* in this region of the spectrum, this also indicates that the companion is probably a late F. Taking the Guide Star Catalogue magnitude of  $V = 13.29$ , the system lies between 755 (F7V) and 870 (F6V) pc away, although if we consider possible errors on the GSC magnitude ( $\pm 0.3 \text{ mag}$ ), and that the absolute magnitudes may be in error by up to  $\pm 0.5 \text{ mag}$ , the system could be as close as 520 pc or as distant as 1250 pc.

Is this a true binary or a chance alignment? Given that the source of the EUV radiation is almost certainly the hot white dwarf, we can calculate the probability of a random 13th magnitude star also falling within the *IUE* aperture. According to Allen (1973), the number of stars per square degree at the Galactic latitude of RE J0500–364, brighter than  $V = 13$ , is 87.1. The *IUE* large aperture is a  $10 \times 20 \text{ arcsec}^2$  oval; hence we calculate a probability of  $\approx 1/250$  of a chance alignment. (Note that the point spread function of the stellar image of the 13th magnitude star on the Digitized Sky Survey plate is  $\sim 15 \text{ arcsec}$  in radius; see Fig. 1).

Assuming this is indeed a true binary, then we believe RE J0500–364 to be one of the most distant white dwarfs, if not the most distant, to be identified in the EUV surveys (for a comparison, see the distance estimates for the white dwarfs detected by the *ROSAT* PSPC by Fleming et al. 1996, and those detected by *EUV* by Vennes et al. 1997c).




**Figure 1.** Field of the *ROSAT* WFC EUV source RE J0500–364 ( $6 \times 6 \text{ arcmin}^2$ ). The circles are the 90 per cent error boxes, centred on the source coordinates, of (in order of decreasing size) *EUV*, the WFC, and the *ROSAT* HRI. The WFC error box is centred on coordinates (J2000)  $05^{\text{h}} 00^{\text{m}} 03^{\text{s}}.9 -36^{\circ} 24' 02''$ . The DA+F6/7V binary lies near the centre of each error box. The arrows indicate other stars examined by Mason et al. (1995) and ourselves during optical searches for the EUV and X-ray counterpart. None of these objects was particularly remarkable.



**Figure 2.** Optical spectrum of RE J0500–364 obtained with the Steward Observatory 2.3-m telescope on Kitt Peak. Using this spectrum, we classify the star as F6/7V.

A model fit to the *IUE* spectrum at  $\log g = 7.25$  gives  $T = 38\,260 \text{ K}$  and  $M = 0.40 M_{\odot}$ , corresponding to a distance of 830 pc. A fit at  $\log g = 8.0$  corresponds to the minimum distance estimate to the system, but gives a higher mass,  $M = 0.68 M_{\odot}$  and higher temperature,  $T = 45\,000 \text{ K}$  (and brings the star into the regime where its atmosphere may be contaminated by elements heavier than He). The inferred  $V$  magnitude from these models is 17.9, making this one of the faintest white dwarfs to be detected by *ROSAT*. The faintest, RE J0616–649 ( $V = 18.4$ ), is a rare magnetic DA (Jordan 1997), but there are few other hot white dwarfs detected fainter than 17th magnitude.




**Figure 3.** Far-UV IUE SWP spectrum of RE J0500–364 (SWP56338), obtained in December 1995 (8-h exposure), clearly showing the white dwarf companion. Also shown is a pure H model atmosphere fit for  $\log g = 7.25$ ,  $T = 38\,260$  K and  $M = 0.40 M_{\odot}$ .

This source is not included in the *ROSAT* PSPC All-Sky Survey Bright Source Catalogue (Voges et al. 1996), but it has been observed and detected in X-rays during a pointed observation by the *ROSAT* High Resolution Imager (HRI). The target was observed as part of programme to identify possible isolated neutron star candidates in the *ROSAT* surveys (PI Wang). Using the Point Source Search programme (PSS) within the Starlink ASTERIX X-ray analysis package (see Pye et al. 1995) to analyse the HRI image, the position of the X-ray source was found to be coincident with the WFC and *EUVE* detections (Fig. 1), and has a flux of  $27.4 \pm 3.9$  counts per  $1000 \text{ s}^{-1}$ . The source shows no evidence for time variability.

Although the HRI has very limited spectral response (the spectral properties of the HRI are known to vary with detector position and time in an ill-defined manner), it is possible to obtain a crude hardness ratio. In this case, the HRI data can then be used to test whether there is any hard X-ray emission from the F6/7V companion star, since no emission is expected from the white dwarf photosphere above 0.4 keV. Fig. 5 shows the HRI data plotted as a function of pulse height distribution. The exact energy of each channel varies temporarily and spatially, but channels 1–5 are roughly equivalent to the soft (0.1–0.4 keV) PSPC band, and channels 6–16 are equivalent to the hard (0.4–2.4 keV) band. As with the PSPC, a hardness ratio can be defined by  $(\text{Hard} - \text{Soft}) / (\text{Hard} + \text{Soft})$ . In this case, the hardness ratio =  $-1.0$ , confirming, as can be readily seen in Fig. 5, that this is a very soft source. We conclude that all of the EUV and soft X-ray emission originates from the hot white dwarf.

Unfortunately, there is no reliable detector matrix for the HRI, and it did not prove possible to use the X-ray count rate in the



**Figure 4.** Far-UV IUE LWP spectrum of RE J0500–364 (LWP31729). The data are very noisy and have been binned up, but the spectral shape most closely matches stars in the range F5V–G0V, consistent with the optical data.

subsequent analysis. Therefore, since this left only the two *ROSAT* WFC data points to try to constrain the interstellar hydrogen column density and white dwarf atmospheric parameters, we assumed that the white dwarf photosphere was essentially pure hydrogen (a reasonable assumption for  $T < 40\,000$  K, and for many hot DAs  $T < 50\,000$  K), and allowed only the neutral hydrogen column density to vary in the subsequent fitting of the EUV data. We found that the WFC S2 photometric data point is well matched by a model

**Table 8.** Neutral hydrogen column and volume densities in the local interstellar medium.

Name	$l$	$b$	$d$ (pc)	$N_{\text{H}}$ ( $\text{cm}^{-2}$ )	$n_{\text{H}}$ ( $\text{cm}^{-3}$ )	Ref.
$\beta$ CMa	226.1	-14.3	206	$2.2 \times 10^{18} \dagger$	0.0035	Cassinelli et al. 1995
$\epsilon$ CMa	239.8	-11.3	188	$1.2 \times 10^{18} \dagger$	0.0021	Cassinelli et al. 1996
RE J0457-281	229.3	-36.2	90	$9.6 \times 10^{17}$	0.0035	Barstow 1997, private comm.
RE J0503-289	230.7	-34.9	90	$8.2 \times 10^{17}$	0.0030	Barstow 1997, private comm.
RE J0500-364	239.6	-36.2	830	$7.5 \times 10^{18}$	0.0029	this paper


$\dagger$  Upper limit

fit at  $\log g = 7.25$ , although the S1 flux was predicted to be slightly lower than observed. The hydrogen column density given by this model is  $N_{\text{H}} = 7.5 \times 10^{18} \text{ atom cm}^{-2}$ . At a distance of 830 pc, this translates into a neutral hydrogen line-of-sight volume density of only  $0.0029 \text{ atom cm}^{-3}$ , well below the average local volume density within  $\sim 80$  pc of the Sun of  $0.05 \text{ atom cm}^{-3}$  (Warwick et al. 1993). If the system is closer (530 pc) and the  $\log g = 8.0$  model is applied, then the H column density is higher ( $N_{\text{H}} = 2.4 \times 10^{19} \text{ atom cm}^{-2}$ ) and the line-of-sight volume density is also higher at  $0.0146 \text{ atom cm}^{-3}$ . Again, however, this is well below the average local volume density.

Notably, this system (galactic coordinates  $l = 239^\circ.6$ ,  $b = -37^\circ.2$ ) lies in a similar direction to the known exceptionally low column densities towards the B1 giant  $\beta$  CMa ( $l = 226^\circ$ ,  $b = -14^\circ$ ; Welsh 1991), and in particular to two other *ROSAT*-discovered hot white dwarfs, RE J0457-281 and RE J0503-289 ( $l = 229^\circ.3$ ,  $b = -36^\circ.2$ ; Barstow et al. 1993).

At a distance of  $\approx 200$  pc,  $\beta$  CMa is known to exist in a rarified ‘interstellar tunnel’ of very low neutral gas density, which is itself an extension of the region surrounding the Sun called the Local Bubble (Welsh 1991). The features of the Bubble were first mapped out by Frisch & York (1983), and Welsh speculates that the  $\beta$  CMa tunnel may extend for *at least* 300 pc in that direction away from the Sun. Welsh also estimates the tunnel to be about 50 pc in diameter; the white dwarfs RE J0457-281 and RE J0503-289 (which are much closer at  $\approx 90$  pc) may then possibly exist in a southward extension of this tunnel, or more likely lie in the foreground, within the Local Bubble itself.

Table 8 details the known column densities and volume densities



**Figure 5.** Pulse height distribution of the X-rays from RE J0500-364 detected by the *ROSAT* HRI. Although the exact energies of each channel are known to vary spatially and temporally, it can be seen that this is a soft source, and there are no hard counts.

to these three stars, and also to  $\epsilon$  CMa (B2 II,  $l = 239^\circ.8$ ,  $b = -11^\circ.3$ ), the brightest EUV source in the sky (Cassinelli et al. 1995), which lies in a similar direction. The line-of-sight neutral hydrogen volume density we measure for REJ0500-364 compares favourably with the average volume density to these four stars ( $0.0030 \text{ atom cm}^{-3}$ ). This suggests that RE J0500-364 might also lie within the  $\beta$  CMa extension of the Local Bubble.

If RE J0500-364 really does lie as far as  $\sim 500$  pc or more away, then it presents a possible lower limit to the size of any neutral gas-free corridor stretching away from the Local Bubble in that direction. The region is bounded on three sides by several OB associations: the Orion nebula (450 pc away), the CMa OB1 association (800 pc distant), and the Gum nebula (290 pc away). Welsh (1991) hypothesizes that this tunnel may have been evacuated by a number of supernova explosions in the last few  $\times 10^5$  yr. The injection of driven, heated, rarefied gas into an older ( $\sim 10^7$  yr old) low-density cavity in the local interstellar medium would produce the large region of very low-density neutral gas that we now see.


### 5.1.2 HD 27483

The Hyades system HD 27483 consists of two active F6V stars orbiting each other with a period of 3.05 d. Although an EUV source was detected originating from the direction of this system in the *ROSAT* WFC all-sky survey in 1990, the hot white dwarf component was identified independently and serendipitously by Böhm-Vitense (1993) during an *IUE* SWP observation as part of a survey of Hyades F stars (Böhm-Vitense 1995). Böhm-Vitense (1993) derived atmospheric parameters for the white dwarf of  $23000 \pm 1000$  K and  $M = 0.6 M_{\odot}$ . However, in the analysis of the white dwarf spectrum the author utilized the unblanketed models of Wesemael et al. (1980), assuming  $\log g = 8.0$ , to fit the far-UV spectrum at just two points. Since the white dwarf might be hot enough to be contributing to the observed EUV flux, it could be argued that the system should be included on the list of EUV-detected hidden white dwarfs, and thus we have decided to re-analyse the far-UV and *ROSAT* data here, in the same manner as the other recently discovered systems.

The *ROSAT* WFC source RE J0420+138, associated with HD 27483, was listed in the original Bright Source Catalogue (Pounds et al. 1993), with a count rate of  $15.0 \pm 4.0 \text{ count ks}^{-1}$  (it was not detected in S2), but was not subsequently detected in the reprocessed 2RE survey (Pye et al. 1995). The significance of the S1 detection in the 2RE survey was 4.2; sources had to exceed a significance of 5.5 in a combination of both bands to be included in the catalogue. Even so, at  $T \approx 23000$  K the white dwarf may be contributing to this small EUV flux, despite the fact that the two F star companions are known to be active themselves.

There is significant contamination in the *IUE* SWP spectrum (see Fig. 6) at the long-wavelength end from the two F6V star





**Figure 6.** Far-UV IUE SWP spectrum of HD 27483 (SWP45940). The hot white dwarf can clearly be seen emerging from the glare of the two F6V companions shortwards of  $\sim 1700$  Å. Also shown is a pure H model atmosphere fit for  $\log g = 8.5$ ,  $T = 22\,000$  K and  $M = 0.94 M_{\odot}$ .

companions, but this falls to zero by  $1600$  Å. We were therefore able to use the continuum flux up to this point. Böhm-Vitense (1993) estimated a distance of  $47.6$  pc to this system. We can further constrain this figure with the recently published *Hipparcos* parallaxes (ESA 1997), where the measured value for HD 27483 is  $21.8 \pm 0.85$  mas, corresponding to a distance of  $45.9 + 1.80/-1.75$  pc. The spectral model which best matches this distance has  $\log g = 8.5$  and  $T = 22\,000$  K, and the corresponding stellar mass is  $M = 0.94 M_{\odot}$ . The white dwarf age is then  $1.4 \times 10^8$  yr, in comparison with the age of the Hyades cluster,  $\sim 7 \times 10^8$  yr (Böhm-Vitense 1993). Given that the white dwarf might have a higher mass than is the average for these stars ( $\approx 0.6 M_{\odot}$ ; Marsh et al. 1997), we can estimate its progenitor mass and place a possible lower limit on the maximum mass for white dwarf progenitor stars in the Hyades cluster. From Wood (1992):

$$M_{\text{WD}} = A \exp(B \times M_{\text{MS}}),$$

where  $A = 0.49462 M_{\odot}$  and  $B = 0.09468 M_{\odot}^{-1}$ . We find, for  $M_{\text{WD}} = 0.94 M_{\odot}$ ,  $M_{\text{MS}} = 6.7 M_{\odot}$ .

The main-sequence lifetime of a  $6.7 M_{\odot}$  star is in fact significantly shorter than  $560$  Myr (the difference between the Hyades age and the white dwarf cooling age; e.g. Schaller et al. 1992). This suggests that the white dwarf is, in reality, probably lower in mass than  $0.94 M_{\odot}$ . However, in order to unambiguously and tightly constrain the fundamental parameters of this star, we will need to obtain a spectrum of the Lyman series with an instrument such as the forthcoming *FUSE* mission (see also Section 6).

The WFC source is coincident with a *ROSAT* PSPC X-ray source, with a total count rate of  $130.6 \pm 19.8$  count  $\text{ks}^{-1}$ , including a detection in the upper band. This confirms that at least one of the

two F6V companion stars is active, as the hard X-rays could not have originated from the white dwarf. The *ROSAT* data points cannot be matched with any of the white dwarf models (which assume a homogeneous atmospheric mixture of H and He). This implies that the active star(s) must be providing a significant fraction of the EUV flux, since little or no heavy-element contamination is expected in the white dwarf photosphere in this cool temperature regime. It is even possible that there is no flux at all from the white dwarf at these wavelengths. The contribution of the white dwarf to the S1 count rate can, however, be estimated. Another hot WD+MS binary, V471 Tauri, is detected by *ROSAT* in the Hyades cluster (Barstow et al. 1992). After subtracting the contribution from the active K2V companion, Marsh et al. (1997) use the WFC count rates to estimate the H column density to this system ( $8.52 \times 10^{18}$  atom  $\text{cm}^{-2}$ ). Adopting the same column density to HD 27483, assuming a pure H atmosphere, and using the parameters derived from the  $\log g = 8.5$  model, the white dwarf is predicted to contribute  $5.4$  count  $\text{ks}^{-1}$  to the S1 flux (i.e.,  $\approx 1/3$  of the  $15$  count  $\text{ks}^{-1}$  detected). How is this count rate affected by uncertainties in the H column density? In fact, from *EUV* spectra, Dupuis et al. (1997) derived a much lower H column density to V471 Tauri of  $1.5 \times 10^{18}$ . Using this value, in the  $\log g = 8.5$  model, the white dwarf contributes  $7.6$  count  $\text{ks}^{-1}$  to the S1 flux (i.e.,  $\approx 1/2$  of the observed flux). This hot degenerate companion to HD 27483 could itself, then, be regarded as a real EUV source. At  $T \approx 22\,000$  K, this would make it one of the coolest white dwarfs to be detected in the EUV surveys.

The combined X-ray luminosity of the two F6V stars in the HD 27483 system can also be estimated, by subtracting the contribution of the white dwarf to the *ROSAT* PSPC lower band flux. The PSPC

count rates are  $98 \pm 14$  count  $\text{ks}^{-1}$  in the softer 0.1–0.4 keV band, and  $33 \pm 9$  count  $\text{ks}^{-1}$  in the harder 0.4–2.4 keV band (Voges et al. 1996). In the  $\log g = 8.5$  model, assuming a column density of  $8.52 \times 10^{18}$  atom  $\text{cm}^{-2}$ , the white dwarf flux in the 0.1–0.4 keV band is found to be 36.0 count  $\text{ks}^{-1}$ . Eliminating this from the total PSPC lower band rate and following the method detailed by Fleming et al. (1995), we find  $L_x = 1.7 \times 10^{29}$  erg  $\text{s}^{-1}$ , and  $L_x/L_{\text{bol}} = 7.8 \times 10^{-6}$ .

## 5.2 Non-detections and active stars

### 5.2.1 BD+49°646 (2RE J0222+50)

This unclassified star was only observed by the SWP camera (SWP56333), and there was no flux visible above the background. If BD+49°646 is a G or K star, then we probably would not detect it in this waveband anyway. No emission features are visible in the UV spectrum, but the EUV source is coincident with a *ROSAT* PSPC hard X-ray source, and the possibility must remain that BD+49°646 is coronally active. Alternatively, the EUV/X-ray source might be another object in the field, or it is possible that the target was missed altogether in the *IUE* observation.

### 5.2.2 2RE J0232–02

As with the WD+MS binary RE J0500–364 (discussed above), the field of this WFC source was originally observed in 1995 with the 2.3-m Steward Observatory telescope at Kitt Peak, as part of a programme to try to identify the remaining optical counterparts to unknown EUV sources in the *ROSAT* WFC catalogues. In the absence of any plausible EUV source, the 15th magnitude G-type central star in the error box may be hiding a hot white dwarf


companion. The far-UV spectra obtained with *IUE* (SWP56272 and LWP31800) were very noisy and showed no evidence for a hot white dwarf. There was some flux above the background longwards of  $\sim 2700$  Å in the LWP spectrum, which may have been due to a G star, but it is possible that the target was missed altogether, and this flux was due to scattered solar light which effects the LWP camera sporadically. At  $V = 14.8$  this star is unlikely to be the source of the EUV flux: a 15th magnitude main-sequence mid-G star would require  $L_{\text{EUV}}/L_{\text{bol}} \sim 0.1$  to produce the count rate seen in the WFC S2 filter (45 count  $\text{ks}^{-1}$ ), far in excess of the saturation level for coronal emission ( $L_{\text{EUV}}/L_{\text{bol}} \sim 10^{-3}$ ; Mathioudakis et al. 1995).

### 5.2.3 SAO 150508 (2RE J0530–19)


Prior to the publication of the 2RE catalogue (Pye et al. 1995), the 9th magnitude F6V star SAO 150508 was not thought to be active, although there are no published optical observations which might offer evidence one way or another. The *IUE* spectra (SWP 56195 and LWP31700; Fig. 7) show no evidence for a hot white dwarf companion. SAO 150508 is, however, coincident with a *ROSAT* PSPC X-ray source. Therefore, estimates of the X-ray and EUV luminosities, assuming that SAO 150508 is active and the true source of the EUV and X-ray flux, are presented in Table 7.

### 5.2.4 HD 36869 (2RE J0534–15)


There is no evidence in the literature that the 8th magnitude G2V star HD 36869 is active. The *IUE* spectra (SWP56169 and LWP31701; Fig. 8) also show no evidence for activity, although the star is coincident with a *ROSAT* PSPC source. Given that 8th magnitude stars are comparatively rare, it is still possible that



**Figure 7.** Far-UV *IUE* spectrum of SAO 150508 (G5V, SWP56195+LWP31700). Clearly, there is no white dwarf companion to this star.



**Figure 8.** Far-UV *IUE* spectrum of HD 36869 (G2V, SWP56169+LWP31701). No white dwarf companion is visible in the short wavelength spectrum.



**Figure 9.** Far-UV *IUE* spectrum of Gl 216B (K2V, SWP56194+LWP31699). Mg II 2798 Å is visible in emission. The feature at ~1720 Å (inset) is probably due to a cosmic ray hit.

HD 36869 is coronally active, and thus we provide estimates of the X-ray and EUV luminosities in Table 7.

### 5.2.5 Gl 216B (2RE J0544–22)

Gl 216B (K2V) is part of a nearby ( $\approx 8$  pc) triple system, and was chosen as a candidate white dwarf binary on the basis of the S2/S1 count rate ratio. The *IUE* spectra (SWP56194 and LWP31699; Fig. 9) show no evidence for a hot white dwarf. However, Mg II 2798 Å is visible in emission (with a line flux of  $4.2 \pm 0.7 \times 10^{-12}$  erg cm $^{-2}$  s $^{-1}$  above the continuum). The emission feature in the SWP spectrum at  $\sim 1720$  Å is probably spurious, since it does not coincide with any commonly seen line. From observations made in the optical, Cayrel de Strobel et al. (1989) concluded that this is a young, active star. If it is the only source of the EUV flux, then we determine  $L_{\text{EUV}}/L_{\text{bol}} = 2.78 \times 10^{-5}$ .

Active stars have a characteristic EUV to Mg II flux ratio. For example, Jewell (1993) shows that the 0.05–0.2 keV EUV flux is  $(1-10) \times$  the Mg II flux. For Gl 216B, however, the EUV to Mg II flux ratio is only  $\approx 0.8$ .

Schmitt et al. (1990) observed the entire Gl 216 system in an *Einstein* HRI pointing, and found that the nearby F7V star Gl 216A was 8 times brighter in X-rays than Gl 216B. Thus Hodgkin & Pye (1994) concluded that all of the EUV radiation in fact comes from Gl 216A. However, we only targeted Gl 216B with *IUE*, since this is the object associated with the EUV source in the 2RE catalogue, and at the time of the observation we were unaware of Hodgkin & Pye's conclusion. The low EUV to Mg II flux ratio for Gl 216B does, however, support these earlier conclusions that the major source of the EUV radiation is actually Gl 216A.


Estimates of the X-ray and EUV luminosities are given in Table 7 assuming that (a) all the flux comes from Gl 216A, and (b) all the flux comes from Gl 216B.

### 5.2.6 HR 2225 (HD 43162, RE J0613–23)


This G5V star was not known to be active prior to the *ROSAT* survey, but subsequently it has been studied in detail by Jeffries & Jewell (1993), and is almost certainly the EUV source. It is also an X-ray source, and measurements of the X-ray and EUV luminosities are given in Table 7. No obvious emission features are visible in the *IUE* LWP and SWP spectra (Fig. 10); the feature longwards of 1800 Å in the SWP spectrum is probably spurious, as there is no commonly seen line at this wavelength.

### 5.2.7 HD 295290 (2RE J0640–03)


There are no references in the literature to this being an active star. However, Mg II is clearly seen in emission at 2800 Å in the *IUE* LWP spectrum (Fig. 11), and there is a suggestion of C IV in emission at 1550 Å in the short-wavelength region. Measurements of the X-ray and EUV luminosities are given in Table 7 using the G0 classification given by SIMBAD, and assuming the star is on the main sequence. The EUV to Mg II flux ratio ( $\approx 8.0$ ) strongly suggests that this star is active and the true source of the EUV radiation. Note that the ratios  $L_{\text{EUV}}/L_{\text{bol}}$  and  $L_{\text{x}}/L_{\text{bol}}$  are significantly larger than for any of the other stars in this sample, approaching the saturated level for coronal emission ( $\sim 10^{-3}$ ). This suggests that the star may be rapidly rotating.



**Figure 10.** Far-UV *IUE* spectrum of HR 2225 (G5V, SWP56206+LWP31715). The emission feature at  $\sim 1800$  Å (inset) is probably spurious.



**Figure 11.** Far-UV *IUE* spectrum of HD 295290 (G0V, SWP56211+LWP31726). Mg II is clearly seen in emission at 2798 Å. C IV 1549 Å emission is also suggested in the SWP spectrum (inset).



**Figure 12.** Far-UV *IUE* spectrum of HD 54402 (K0, SWP56193+LWP31698). The feature at  $\sim 1800$  Å is probably due to a cosmic ray hit.

5.2.8 *HD 54402 (2RE J0710+45)*

No references are given in the literature to this being an active star. There is clearly no white dwarf visible in the *IUE* SWP spectrum (Fig. 12), and the emission line at  $\sim 1800$  Å is probably spurious, perhaps due to a cosmic ray hit. The EUV source is not coincident with an X-ray source. The star needs to be examined optically to search for any evidence of chromospheric activity.

5.2.9 *SAO 135659 (2RE J0813-07)*

Again, this star was not known to be active prior to the *ROSAT* survey. The *IUE* LWP spectrum (Fig. 13) reveals Mg II in emission at 2800 Å. Measurements of the EUV and X-ray luminosities (this star is also a PSPC source) are given in Table 7, assuming the K0 spectral type given by SIMBAD, and that the star is on the main sequence. The EUV to Mg II flux ratio ( $\approx 8.25$ ) strongly suggests that this star is active and the true source of the EUV radiation.

5.2.10 *RE J0823-252/HD 70907*

With S1 and S2 count rates of  $52 \pm 7$  and  $83 \pm 9$  count  $\text{ks}^{-1}$ , this is a relatively bright EUV source in comparison with most of the targets in this paper. The soft X-ray and EUV photometric colours are also characteristic of a hot white dwarf, and it is not detected in the PSPC hard band. Therefore, it was selected as a potential hidden white dwarf binary.


The  $V = 8.8$  star in the centre of the field, HD 70907 (F3IV/V), was observed in both the *IUE* SWP and LWP cameras (Fig. 14). There is no evidence for a white dwarf companion or emission

features indicative of an active star. A nearby  $V \approx 11$  star was also observed, and again there was no evidence for a white dwarf, although, in the absence of any flux in the LWP camera that could be attributed to a stellar source, it seems possible that the star was not in the LWP slit. Mason et al. (1995) report that this fainter object is indeed active, although they give no indication of the size of any emission features seen in the optical. They also do not give a spectral type for this star, and thus we have not been able to determine the EUV and X-ray luminosities. Whether this star is active enough to be the true EUV source remains unclear, and the suspicion remains that there is indeed an unresolved hot white dwarf hiding in this field.


5.2.11 *HR 4646 (2RE J1212+77)*

It is highly unusual to detect an A star in EUV or X-ray surveys (Fleming et al. 1991). Observations by the *Einstein* and *EXOSAT* observatories failed to find any convincing detections other than the nearby quadruple A star system Castor (Pallavicini et al. 1990). Therefore HR 4646, an Am star coincident with *ROSAT* WFC 2RE and PSPC sources, was selected as a potential hidden white dwarf binary. It should be noted that Am stars do not possess significant magnetic fields and they are slow rotators, but they almost always appear to lie in close binary systems (Abt 1961), and indeed Margoni, Munari & Stagni (1992) found that HR 4646 is a spectroscopic binary with a period of 1.27 d.


The *IUE* SWP spectrum (Fig. 15) shows no evidence for a hot white dwarf companion. It is extremely likely, therefore, that HR 4646 has a coronally active cooler companion (F5 or later).



**Figure 13.** Far-UV *IUE* spectrum of SAO135659 (K0, SWP56273+LWP31801). Mg II is visible in emission at 2798 Å.



**Figure 14.** Far-UV *IUE* spectrum of HD 70907 (F3IV/V, SWP56344+LWP31836). There is no evidence for activity.



**Figure 15.** Far-UV *IUE* SWP spectrum of HR 4646 (A5m, SWP55658). Clearly, there is no white dwarf companion to this star.

## 6 SUMMARY AND CONCLUSIONS

A search for unresolved white dwarfs in binary systems with optically brighter normal stellar companions has been conducted with *IUE* during its final year of operation. Targets were chosen among the fainter and still unidentified EUV sources in the *ROSAT* 2RE catalogue (Pye et al. 1995). One new system was discovered (RE J0500–364, DA+F6/7V), which appears to lie in a direction of low interstellar neutral hydrogen volume density. If this star is really as distant as  $\sim 500$ – $1000$  pc, then it may represent a lower limit to the size of the  $\beta$  CMa tunnel of low-density neutral gas stretching away from the Local Bubble in that direction.

Including the independently identified HD 27483 (DA+F6V; Böhm-Vitense 1993), this new discovery brings the total number of such systems found in the *ROSAT* and *EUVE* surveys to 19. These two satellites have, therefore, been very successful in helping us to identify these kinds of binary systems, which have never been seen optically and could only have been identified through such satellite surveys. However, these stars still represent <20 per cent of the hot white dwarfs identified in the EUV.

The demise of *IUE*, which suffered a gyro failure in 1996 February, limiting it to observe only targets with bright guide stars (the mission was finally terminated in 1996 September), effectively cut short our programme. The loss of *IUE* means that this particular method of searching for these important systems is no longer available. Although *HST* can observe the same wavelength region, it is of course much harder to obtain the time required for this kind of search. It is likely, therefore, that very few new examples of these systems will be discovered in the near future. The results presented in this paper seem to suggest that this search, using the EUV catalogues as a basis from which to identify potential systems, has been fairly exhaustive (we have identified only one new system from 13 targets).

However, the identification by Vennes et al. (1997b) of a hot white dwarf companion to a previously catalogued EUV-bright active star, RE J0702+129 (K0Ve), instead suggests that, in fact, some systems may have been completely overlooked. RE J0702+129 was classified as active by Mason et al. (1995) in their optical follow-up programme to identify the EUV sources found in the *ROSAT* WFC survey, and was not, therefore, included on any of our target lists for the *IUE* search. This raises the question of how many of the >200 ‘active’ stars in the WFC and *EUVE* catalogues really are the source, or at least the only source, of the EUV radiation. Until each star has been observed and analysed in detail, the suspicion remains that there are more of these binary systems in the survey waiting to be identified.

For example, the *ROSAT* WFC catalogue includes  $\approx 120$  isolated white dwarfs, and  $\approx 60$  in some kind of binary, e.g., classic Sirius-type systems, CVs, non-interacting DA+dM systems, and visual binaries. Conservatively, then, we might assume that we have already identified the majority of the white dwarf binaries to be found in the survey. However, if as many as 80 per cent of stars reside in binary or multiple systems, another 30 might be awaiting discovery. These could include double-degenerate systems as well as further examples of Sirius-type systems (e.g., the apparently isolated DA RE J0512–007 has a mass,  $M = 0.38 M_{\odot}$ , too low for it to be the result of single-star evolution, and may have a degenerate companion).

What is really needed to try to find more Sirius-type binaries is an all-sky UV survey. None has been undertaken since the TD-1 survey of 1972/73, which originally appeared to have only detected Sirius B among these systems. Subsequently, though, Landsman, Simon

& Bergeron (1996) found white dwarf companions to 56 Persei and HR 3643 as a result of the UV excess detected by TD-1. The recently approved *GALEX* mission (Bianchi 1998), which will survey the sky at UV wavelengths and follow-up some targets spectroscopically, may reveal many more of these binaries.

In the meantime, follow-up observations of these 19 new EUV-bright systems are required. In particular, it is important to determine whether these systems are wide, or close enough that they must have undergone common-envelope evolution. This information will help to place constraints on theoretical models of binary evolution (e.g. de Kool & Ritter 1993). Detailed studies of the normal stellar companions may also reveal evidence of past interaction (e.g. Jeffries & Stevens 1996), and stars with possible abundance anomalies that may be the progenitors of the barium and carbon giants (e.g. Jeffries & Smalley 1996).

In addition, the forthcoming *FUSE* mission (*Far Ultraviolet Spectroscopic Explorer*) will, for the first time, allow us to unambiguously determine the temperatures and surface gravities of the white dwarfs in these systems (and hence their masses and radii), through modelling of the Lyman absorption series down to 912 Å.

## ACKNOWLEDGMENTS

MRB and MAB acknowledge the support of PPARC. We thank Detlev Koester (Kiel) for the use of his white dwarf model atmosphere grids, and Jim Collins (Arizona) and Nigel Bannister (Leicester) for their help in obtaining and reducing the optical spectrum of RE J0500–364. MRB thanks the staff at *IUE* Vilspa for their help and co-operation in carrying out this programme. We also thank the referee, Rob Jeffries, for useful suggestions to improve this paper. This research has made use of the SIMBAD data base operated at CDS, Strasbourg, France, and data obtained from the Leicester Database and Archive Service (LEDAS) at the Department of Physics and Astronomy, Leicester University, UK.

## REFERENCES

- Abt H. A., 1961, *ApJS*, 6, 37
- Allen C., 1973, *Astrophysical Quantities*. Athlone Press, London
- Barstow M. A., Schmitt J. H. M. M., Clemens J. C., Pye J. P., Denby M., Harris A. W., Pankiewicz G. S., 1992, *MNRAS*, 255, 369
- Barstow M. A. et al., 1993, *Advances in Space Research*, 131, 281
- Barstow M. A. et al., 1994, *MNRAS*, 270, 499
- Bianchi L., 1998, *Ultraviolet Astrophysics – Beyond the IUE Final Archive*, ESA Publication SP–413, p. 797
- Bohlin R. C., Grillmair C. J., 1988, *ApJS*, 66, 209
- Böhm-Vitense E., 1980, *ApJ*, 239, L79
- Böhm-Vitense E., 1993, *AJ*, 106, 1113
- Böhm-Vitense E., 1995, *AJ*, 110, 228
- Bowyer S. et al., 1994, *ApJS* 93, 569
- Bowyer S. et al., 1996, *ApJS*, 102, 129
- Burleigh M. R., Barstow M. A., 1997, in Isern J., Hernanz M., Garcia-Berro E., eds, *White Dwarfs*. Kluwer, Dordrecht, p. 329
- Burleigh M. R., Barstow M. A., 1998, *MNRAS*, 295, L15
- Burleigh M. R., Barstow M. A., Fleming T. A., 1997, *MNRAS*, 287, 381 (Paper I)
- Cassinelli J. P. et al., 1995, *ApJ*, 438, 932
- Cassinelli J. P. et al., 1996, *ApJ*, 460, 949
- Cayrel de Strobel G., Lebreton Y., Perrin M.-N., Cayrel R., 1989, *A&A*, 225, 369
- de Kool M., Ritter H., 1993, *A&A*, 267, 397
- Dupuis J., Vennes S., Chayer P., Cully S., Rodriguez-Bell T., 1997, in Isern J., Hernanz M., Garcia-Berro E., eds, *White Dwarfs*. Kluwer, Dordrecht, p. 277



- ESA, 1997, The *Hipparcos* Catalogue, ESA SP–1200
- Fleming T. A., Liebert J., Green R. G., 1986, *ApJ*, 308, 176
- Fleming T. A., Schmitt J. H. M. M., Barstow M. A., Mittaz J. P. D., 1991, *A&A*, 246, L47
- Fleming T. A. et al., 1995, *ApJS*, 99, 701
- Fleming T. A., Snowden S. L., Pfefferman E., Briel U., Greiner J., 1996, *A&A*, 316, 147
- Frisch P. C., York D. G., 1983, *ApJ*, 271, L59
- Garhart M., 1997, *IUE Newsletter* (GSFC) Vol. 5, No. 5
- Hodgkin S. T., Pye J. P., 1994, *MNRAS*, 267, 840
- Hodgkin S. T., Barstow M. A., Fleming T. A., Monier R., Pye J. P., 1993, *MNRAS*, 263, 229
- Jacoby J. H., Hunter D. A., Christian C. A., 1984, *ApJS*, 56, 257
- Jeffries R. D., 1995, *MNRAS*, 273, 559
- Jeffries R. D., Jewell S. J., 1993, *MNRAS*, 264, 106
- Jeffries R. D., Smalley B., 1996, *A&A*, 315, L19
- Jeffries R. D., Stevens I. R., 1996, *MNRAS*, 279, 180
- Jeffries R. D., Burleigh M. R., Robb R. M., 1996, *A&A*, 305, L45
- Jewell S. J., 1993, PhD thesis, Univ. Birmingham
- Johnson H. R., Ake T. B., 1986, ESA SP-263, 395
- Jordan S., 1997, in Isern J., Hernanz M., Garcia-Berro E., eds, *White Dwarfs*. Kluwer, Dordrecht, p. 397
- Koester D., 1991, in Michaud G., Tutukov A., eds, *Proc. IAU Symp. 145, Evolution of Stars: The Photospheric Abundance Connection*. Kluwer, Dordrecht, p. 435
- Landsman W., Simon T., Bergeron P., 1996, *PASP*, 108, 250
- Margoni R., Munari U., Stagni R., 1992, *A&AS*, 93, 545
- Marsh M. C. et al., 1997, *MNRAS*, 286, 369
- Mason K. O. et al., 1995, *MNRAS*, 274, 1194
- Mathioudakis M., Fruscione A., Drake J. J., McDonald K., Bowyer S., Malina R. F., 1995, *A&A*, 300, 775
- Pallavicini R., Tagliaferri G., Pollock A. M. T., Schmitt J. H. M. M., Rosso C., 1990, *A&A*, 227, 483
- Pounds K. A. et al., 1993, *MNRAS*, 260, 77
- Pye J. P. et al., 1995, *MNRAS*, 274, 1165
- Schafer R. A. et al., 1991, ESA TM-09
- Schaller G., Schaerer D., Meynet G., Maeder A., 1992, *A&AS*, 96, 269
- Schindler M., Stencel R. E., Linsky J. L., Basri G. S., Helfand D. J., 1982, *ApJ*, 263, 269
- Schmitt J. H. M. M., Collura A., Sciortino S., Vaiana G. S., Harnden F. R., Rosner R., 1990, *ApJ*, 365, 704
- Shipman H. L., Geczi J., 1989, in Wegner G., ed., *White Dwarfs*. Springer-Verlag, Berlin, p. 134
- Vennes S., Mathioudakis M., Doyle J. G., Thorstensen J. R., Byrne P. B., 1995, *A&A*, 299, L29
- Vennes S., Berghoefer T., Christian D. J., 1997a, *ApJ*, 491, L85
- Vennes S., Christian D. J., Mathioudakis M., Doyle J. G., 1997b, *A&A*, 318, L9
- Vennes S., Thejll P. A., Genova Galvan R., Dupuis J., 1997c, *ApJ*, 480, 714
- Voges W. et al., 1996, *IAU Circ.* 6420
- Warwick R. S., Barber C. R., Hodgkin S. T., Pye J. P., 1993, *MNRAS*, 262, 289
- Welsh B. Y., 1991, *ApJ*, 373, 556
- Wesemael F., Auer L. H., van Horn H. M., Savedoff M. P., 1980, *ApJS*, 43, 159
- Wood M. A., 1992, *ApJ*, 386, 539
- Wood M. A., 1995, in Koester D., Werner K., eds, *Proc. of the 9th European Workshop on White Dwarfs*. Springer, Berlin, p. 41
- Wu C.-C. et al., 1992, *IUE Ultraviolet Spectral Atlas of Selected Astronomical Objects*, NASA Reference Publication 1285

This paper has been typeset from a  $\text{T}_{\text{E}}\text{X}/\text{L}^{\text{A}}\text{T}_{\text{E}}\text{X}$  file prepared by the author.